

# Community-Based Primary Palliative Care Community of Practice Series 2

## Palliative Sedation



**Facilitator:** Dr. Haley Draper

**Presenter:** Elisabeth Antifeau and Kevin Wade

**Date:** March 15<sup>th</sup> 2023

# Territorial Honouring



# The Palliative Care ECHO Project

The Palliative Care ECHO Project is a 5-year national initiative to cultivate communities of practice and establish continuous professional development among health care providers across Canada who care for patients with life-limiting illness and their families.

**Stay connected: [www.echopalliative.com](http://www.echopalliative.com)**

The Palliative Care ECHO Project is supported by a financial contribution from Health Canada. The views expressed herein do not necessarily represent the views of Health Canada.



# LEAP Core

- Interprofessional course that focuses on the essential competencies to provide a palliative care approach.
- Taught by local experts who are experienced palliative care clinicians and educators.
- Delivered online or in-person.
- Ideal for any health care professional (e.g., physician, nurse, pharmacist, social worker, etc.) who provides care for patients with life-threatening and progressive life-limiting illnesses.
- Accredited by the CFPC and Royal College.



Learn more about the course and topics covered by visiting

[www.pallium.ca/course/leap-core](http://www.pallium.ca/course/leap-core)

# Objectives of this Series

**After participating in this series, participants will be able to:**

- Augment their primary-level palliative care skills with additional knowledge and expertise related to providing a palliative care approach.
- Connect with and learn from colleagues on how they are providing a palliative care approach.

# Overview of Sessions

Session #	Session Title	Date/ Time
Session 1	Pain: Beyond the Basics	Nov 9, 2022 from 1-2pm ET
Session 2	Communication: Part 1	Nov 23, 2022 from 1-2pm ET
Session 3	Communication: Part 2	Dec.7, 2022 from 1-2pm ET
Session 4	Palliative Care and Substance Use Disorders	Jan 18, 2023 from 1-2pm ET
Session 5	GI Symptoms in Palliative Care	Feb 1, 2023 from 1-2pm ET
Session 6	Delirium	Feb 15, 2023 from 1-2pm ET
Session 7	Spiritual Care and Rituals around Death and Dying	Mar 1, 2023 from 1-2pm ET
Session 8	Palliative Sedation	Mar 15, 2023 from 1-2pm ET
Session 9	Grief and Bereavement: Beyond the Basics	Apr 12, 2023 from 1-2pm ET
Session 10	Practical Tips: Lessons from the Front Line	Apr 26, 2023 from 1-2pm ET

# Welcome & Reminders

- Please introduce yourself in the chat! Let us know what province you are joining us from, your role and your work setting.
- Your microphones are muted. There will be time during this session when you can unmute yourself for questions and discussion.
- You are welcome to use the chat function to ask questions and add comments throughout the session.
- This session is being recorded and will be emailed to registrants within the next week.
- Remember not to disclose any Personal Health Information (PHI) during the session.
- This 1-credit-per-hour Group Learning program has been certified by the College of Family Physicians of Canada for up to **11 Mainpro+** credits.

# Disclosure

Relationship with Financial Sponsors:

## **Pallium Canada**

- Not-for-profit
- Funded by Health Canada



# Disclosure

## **This program has received financial support from:**

- Health Canada in the form of a contribution program.
- Generates funds to support operations and R&D from Pallium Pocketbook sales and course registration Fees.

## **Facilitator/ Presenter:**

- Dr. Nadine Gebara: Nothing to disclose.
- Elisabeth Antifeau: Nothing to disclose.
- Kevin Wade: Chief Medical Officer of Gravitii.care, a startup platform to connect home care patients directly with providers.

# Disclosure

## Mitigating Potential Biases:

- The scientific planning committee had complete independent control over the development of course content.

# Introductions

## Facilitator:

### **Dr. Haley Draper, MD CCFP- PC**

Clinical co-lead of this ECHO series

Palliative Care Physician at Toronto Western Hospital, University Health Network

Family Physician at Gold Standard Health, Annex

## Panelists:

### **Dr. Nadine Gebara, MD CCFP- PC**

Clinical co-lead of this ECHO series

Palliative Care Physician at Toronto Western Hospital, University Health Network

Family Physician at Gold Standard Health, Annex

### **Dr. Roger Ghoche, MDCM CCFP-PC, MTS**

Palliative Care and Rehabilitation Medicine, Mount Sinai Hospital- Montreal

# Introductions

## Panelists (continued):

**Thandi Briggs, RSW MSW**

Care Coordinator, Integrated Palliative Care Program  
Home and Community Care Support Services Toronto Central

**Claudia Brown, RN BSN**

Care Coordinator, Integrated Palliative Care Program  
Home and Community Care Support Services Toronto Central

**Rev. Jennifer Holtslander, SCP-Associate, MRE, BTh**

Spiritual Care Provider

## Support Team

**Holly Finn**

Senior Manager, Program Delivery, Pallium Canada

**Darwin Namata**

Program Delivery Officer, Pallium Canada

# Introductions

## **Presenter:**

### **Elisabeth Antifeau, RN, MScN, CHPCN(C), GNC(C)**

Regional Clinical Nurse Specialist (CNS-C), Palliative End of Life Care  
IH Regional Palliative End of Life Care Program  
Pallium Canada Master Facilitator & Coach, Scientific Consultant

### **Kevin Wade, CD, MD, CCFP (PC)**

Palliative Care Physician, BC Cancer and Island Health, Victoria, BC  
Clinical Assistant Professor, University of British Columbia  
Major, 1 Canadian Field Hospital Det Ottawa


# Clinician Considerations for Palliative Sedation

# Session Learning Objectives

**Upon completing the session, participants will be able to:**

- Identify indications for palliative sedation therapy
- Explain the difference between sedation and analgesia
- Explain pharmacotherapy for palliative sedation therapy

# Indications



- Patient at end of life
  - Usual **prognosis < 1wk**, sometimes extended up to 2 wks after careful evaluation
- Refractory Symptoms
  - Most commonly **confusion/delirium**
  - Less commonly pain, dyspnea, nausea, anxiety
  - “refractory” after discussion of available therapies and their likelihood of efficacy, with patient or SDM. Possible trial of therapy
- Setting Capable of handling monitoring and medication administration
  - Usually **inpatient/hospice**




# Indications – Determining Refractory Symptoms

- Determining whether symptoms are refractory:
  - Always requires **discussion with the patient or SDM**
  - May require **consultation with a subject matter** e.g., palliative care physician, anesthesia/pain, internal medicine
  - May require a **trial of therapy** e.g., typical antipsychotics, CADD pump, epidural



# Sedation vs Analgesia

- Quiet does not equal comfortable.
- Consider sedation as an independent factor from other symptom measures.
  - Mostly using RASS-Pal
  - Pasero scale not appropriate
- Manage symptoms as well as possible, even in the sedated patient.



## Richmond Agitation Sedation Scale – Palliative Version (RASS-PAL)

Score	Term	Description
+4	Combative	Overtly combative, violent, immediate danger to staff, (e.g., throwing items): +/- attempting to get out of bed or chair
+3	Very Agitated	Pulls or removes lines (e.g. IV/SC/Oxygen tubing) or catheter(s); aggressive, +/- attempting to get out of bed or chair
+2	Agitated	Frequent non-purposeful movement, +/- attempting to get out of bed or chair
+1	Restless	Occasional non-purposeful movement, but movements are not aggressive or vigorous
0	Alert and Calm	
-1	Drowsy	Not fully alert but has sustained awakening (eye-opening / eye contact) to voice for 10 seconds or longer.
-2	Light Sedation	Briefly awakens with eye contact to voice for less than 10 seconds
-3	Moderate Sedation <i>(common goal)</i>	Any movement (eye or body) or eye opening to voice, but no eye contact
-4	Deep Sedation	No response to voice but any movement (eye or body) or eye opening to stimulation by light touch
-5	Not rousable	No response to voice or stimulation by light touch

# Pain Assessment In Advanced Dementia (PAIN-AD) Scale

	0	1	2	Score
<b>Breathing</b> Independent of vocalization	Normal	Occasional labored breathing. Short period of hyperventilation.	Noisy labored breathing. Long period of hyperventilation. Cheyne-Stokes respirations.	
<b>Negative Vocalization</b>	None	Occasional moan or groan. Low level speech with a negative or disapproving quality.	Repeated troubled calling out. Loud moaning or groaning. Crying.	
<b>Facial Expression</b>	Smiling or inexpressive	Sad. Frightened. Frown.	Facial grimacing.	
<b>Body Language</b>	Relaxed	Tense. Distressed pacing. Fidgeting.	Rigid. Fists clenched, knees pulled up. Pulling or pushing away. Striking out.	
<b>Consolability</b>	No need to console	Distracted or reassured by voice or touch.	Unable to console, distract or reassure.	
<b>TOTAL</b>				

- Scoring:**
- 1–3 Mild pain *Provide comfort measures (i.e., non-pharmacologic approaches such as repositioning or distraction or a mild analgesic such as acetaminophen)*
  - 4–6 Moderate pain
  - 7–10 Moderate to Severe pain *Pain that warrants stronger analgesia, such as an opioid, as well as comfort measures*

# Pharmacology

- **Neuroleptics**
  - Methotrimeprazine (Nozinan)
  - Haloperidol (Haldol)
- **Benzodiazepines**
  - Midazolam
  - Lorazepam
- **Less common**
  - Barbiturates (e.g., Phenobarbital)
  - Surgical anesthetics (e.g. Propofol, Dexmedetomidine)

# Questions



# Nursing Considerations for Palliative Sedation

# Overview of the Palliative Sedation Assessment Tools for Nursing

## What to Assess

- Level of Sedation
- Level of Pain (non-communicative)
- Level of Respiratory Distress/Dyspnea
- Integration of Assessment and Decision-Making

## Recommended Tool

- RASS-PAL (Richmond Agitation Sedation Scale – Palliative Care)
- PAINAD – Pain Assessment in Advanced Dementia
- RDOS – Respiratory Distress Observation Scale
- Palliative Sedation Monitoring Record with the assessment tools integrated into the record



# Level of Sedation

- Tool permits standardized assessment of level of sedation – as ordered goal (target) and maintenance.
- RASS-PAL:
  - Valid and reliable assessment tool for palliative sedation;
  - 10 Point scale:
  - 0 = Alert and Calm
    - +1 to +4 escalates from restless to combative
    - -1 to -5 descends from drowsy to non-rousable
  - Sedation goal is commonly ordered at -3 (Moderate Sedation)
  - No noxious stimuli
  - Procedure is standardized and part of the tool

# Level of Pain

- Need a tool that permits standardized assessment of pain in non-communicative patients.
- PAINAD (Pain Assessment in Advanced Dementia) Scale:
  - Score 5 areas of described observational data in a 3 point scale (0 – 1 – 2)
    - Breathing
    - Negative Vocalization
    - Facial Expression
    - Body Language
    - Consolability
- Scoring out of 10: Mild Pain (=1-3); Moderate Pain (=4-6) and Mod-Severe Pain (7-10)

# Level of Respiratory Distress/Dyspnea

- Standardized assessment tool for respiratory distress (non-communicative).
- RDOS (Respiratory Distress Observation Scale)
  - Assesses 8 variables in a 3 point scale (0-1-2);
    - Heart Rate/min and Respiratory Rate/min
    - Restlessness (non-purposeful movements)
    - Paradoxical breathing
    - Accessory muscle use
    - Grunting at end-expiration
    - Nasal flaring
    - Look of fear (described)
  - RDOS < 3 = respiratory comfort; >3 = respiratory distress, need for palliation.

# Clinical Decision Making & Documentation

- Quiet does not = Comfort!
- Clinical Decision Making and Problem Solving:
  - Assess PAINAD, RDOS and RASS-PAL Scores each time patient is restless/appears to be more awake.
  - Treat PAINAD and RDOS scores with opiates first before sedating.
  - Reassess PAINAD and RDOS 3-5 minutes later,
    - if still elevated give more opioids;
    - if within normal levels (PAINAD 3 or less; RDOS 3 or less) – give sedation bolus
- Best practice is to use a monitoring record that integrates correct tool use with clinical decision-making and guides decisions re analgesia vs sedation

# Case-Based Discussion



# Case Introduction



**ID:** 67M with new diagnosis glioblastoma. Non-operative. Pending radiation oncology consult. Came to ED for falls/confusion. Increasing delirium and agitation, admitted under hospitalist

**Social situation:** Lives with his wife and one adult son. Two other adult children live away from town

**PMHx:** Meningioma 6yrs ago, resected. Controlled HTN, Impaired Fasting Glucose

Goals of care not established yet

**WHAT IS YOUR APPROACH?**

# Case Cont'd

- Workup of delirium demonstrates no clear reversible cause
- Extensive vasogenic edema surrounding L parietal lobe lesion on MRI
  - No improvement after 48hr trial of dex. Repeat CT shows decreased edema but still delirious/agitated. Occasionally in restraints, family holding him down.  
?partial seizures
- Goals of care conversation by hospitalist and social worker after 10 days – family are exhausted. Request comfort care

# Case – Palliative

- **Where do you start?**
  - **Is this patient at the end of their life?**
  - **Do they have refractory symptoms?**
- **What are the challenges you currently face?**
- **What agent would you choose for sedation?**

**Nobody  
should die in  
restraints!**



# Session Wrap Up

- Please fill out our feedback survey, a link has been added into the chat.
- A recording of this session will be emailed to registrants within the next week.
- We hope to see you again at our next session taking place **April 12, 2023 from 1-2pm ET** on the topic of **Grief and Bereavement: Beyond the Basics.**

# Thank You



Stay Connected  
[www.echopalliative.com](http://www.echopalliative.com)